

Liquid Preparations Available	Formulations	Usual Doses	Total Daily Liquid Amount
Fosamprenavir (Lexiva®)	<ul style="list-style-type: none"> • Available in Spring 2007 • Oral suspension (50mg/mL) in a 225 mL bottle and does not reconstitution • Off-white color with grape, bubblegum or peppermint flavor • Stored at 40-80°F, although refrigeration improves palatability • Shake well before using • Take on an empty stomach 	<ul style="list-style-type: none"> • <i>Naive pts only:</i> 28 mL BID • <i>Naive pts only:</i> 28 mL daily + rTV 2.5 mL daily • FPV 14 mL BID + rTV 1.25 mL BID 	<ul style="list-style-type: none"> • <i>Naive pts only:</i> 56 mL • <i>Naive pts only:</i> FPV 56 mL + rTV 2.5mL = 58.5 mL • FPV 14 mL BID + rTV 1.25 mL = 29.25 mL
Ritonavir (Norvir®)	<ul style="list-style-type: none"> • Oral Solution containing 600 mg/7.5 mL in 240 mL bottle with measuring cup • Refrigeration of oral solution is not required • Patients find preparation unpalatable since it is a castor-oil base containing alcohol and propylene glycol • Peppermint-caramel flavor 	<ul style="list-style-type: none"> • With other PI: 100 (1.25 mL) to 200 (2.5 mL) mg Q12-24H 	<ul style="list-style-type: none"> • 1.25-5 mL
Lopinavir/ritonavir (Kaletra®)	<ul style="list-style-type: none"> • Oral Solution (lopinavir/ritonavir: 400 mg/100 mg per 5 mL) • Equivalent to taking two of the tablets • Comes in bottles of 160 mL • Cotton-candy-flavored liquid containing propylene glycol and is 42.4% alcohol 	<ul style="list-style-type: none"> • 5 mL BID with food • If pt is taking concomitant efavirenz or nevirapine, may consider increasing dose to 6.5 mL BID with food 	<ul style="list-style-type: none"> • 10 mL • <i>Concomitant use with efavirenz or nevirapine, 13 mL</i>
Tipranavir (Aptivus®)	<ul style="list-style-type: none"> • Oral Solution: 100 mg/mL in 95 mL bottle • Contains vitamin E, propylene glycol; butter mint-butter toffee flavor 	<ul style="list-style-type: none"> • 5 mL BID + rTV 2.5mL BID 	<ul style="list-style-type: none"> • 15 mL
Nelfinavir (Viracept®)	<ul style="list-style-type: none"> • Available as an oral powder containing 50 mg nelfinavir base per one gram scoop • Patients should mix correct amount of powder with water or milk and ingest within 6 hours • DO NOT mix with acidic juices or apple sauce as it may result in a bitter taste 	<ul style="list-style-type: none"> • Mix 25 grams with small amount of water or milk BID 	
Atazanavir (Reyataz®)	<ul style="list-style-type: none"> • Capsules can be opened, mixed with 4 oz. of applesauce & ingested immediately prior to a light meal (Bristol-Meyier Squibb) • Pharmacokinetic study documented that bioavailability of atazanavir administered in applesauce was 92% compared to ingestion of two 200 mg capsules, both within 5 minutes of a light meal. Although peak concentration was lower when capsules were opened, the AUCs were similar. 	<ul style="list-style-type: none"> • <i>Naive pt only:</i> 400 mg daily • ATV 300 mg daily + rTV 1.25 mL daily 	